







critically, contributed to the interpretation of data and approval of the submitted and final versions.

## References

- 1) Steib A, Freys G, Lehmann C, et al. Intraoperative blood losses and transfusion requirements during adult liver transplantation remain difficult to predict. *Can J Anaesth* 2001; **48**: 1075-9.
- 2) Rana A, Petrowsky H, Hong JC, et al. Blood transfusion requirement during liver transplantation is an important risk factor for mortality. *J Am Coll Surg* 2013; **216**: 902-7.
- 3) Benson AB, Burton JR, Austin GL, et al. Differential effects of plasma and red blood cell transfusions on acute lung injury and infection risk following liver transplantation. *Liver Transpl* 2011; **17**: 149-58.
- 4) Azevedo LD, Stucchi RS, Ataíde EC, Boin IF. Assessment of causes of early death after twenty years of liver transplantation. *Transplant Proc* 2013; **45**: 1116-8.
- 5) Massicotte L, Lenis S, Thibeault L, et al. Effect of low central venous pressure and phlebotomy on blood product transfusion requirements during liver transplantations. *Liver* 2006; **12**: 117-23.
- 6) Coakley M, Reddy K, Mackie I, et al. Transfusion triggers in orthotopic liver transplantation: a comparison of the thromboelastometry analyzer, the thromboelastogram, and conventional coagulation tests. *J Cardiothorac Vasc Anesth* 2006; **20**: 548-53.
- 7) Spahn DR, Theusinger OM, Hofmann A. Patient blood management is a win-win: a wake-up call. *Br J Anaesth* 2012; **108**: 889-92.
- 8) Cywinski JB, Alster JM, Miller C, et al. Prediction of Intraoperative Transfusion Requirements During Orthotopic Liver Transplantation and the Influence on Postoperative Patient Survival. *Anesth Analg* 2014; **118**: 428-37.
- 9) Escorcesca Ortega AM, Mogollon Jiménez MV, Hinojosa Pérez R, et al. Application of the McCluskey Index to Predict Blood Product Requirements During Liver Transplantation. *Transplant Proc* 2008; **40**: 2981-2.
- 10) Findlay JY, Rettke SR. Poor prediction of blood transfusion requirements in adult liver transplantations from preoperative variables. *J Clin Anesth* 2000; **12**: 319-23.
- 11) Freeman RB Jr, Wiesner RH, Harper A, et al. The new liver allocation system: moving toward evidence-based transplantation policy. *Transpl* 2002; **8**: 851-8.
- 12) Shander A, Van Aken H, Colomina MJ, et al. Patient blood Management in Europe. *Br J Anaesth* 2012; **109**: 55-68.
- 13) Hosmer DW Jr, Lemeshow S, Sturdivant RX, et al. *Applied Logistic Regression*. 3<sup>rd</sup> ed. Hoboken: Wiley; 2013.
- 14) Massicotte L, Sassine MP, Lenis S, et al. Transfusion predictors in liver transplant. *Anesth Analg* 2004; **98**: 1245-51.
- 15) Mangus RS, Kinsella SB, Nobari MM, et al. Predictors of blood product use in orthotopic liver transplantation using the piggyback hepatectomy technique. *Transplant Proc* 2007; **39**: 3207-13.
- 16) Pihusch R, Rank A, Gohring P, et al. Platelet function rather than plasmatic coagulation explains hypercoagulable state in cholestatic liver disease. *J Hepatol* 2002; **37**: 548-55.
- 17) Ben-Ari Z, Panagou M, Patch D, et al. Hypercoagulability in patients with primary biliary cirrhosis and primary sclerosing cholangitis evaluated by thrombelastography. *J Hepatol* 1997; **26**: 554-9.

---

Arrived: 10 August 2015 - Revision accepted: 28 October 2015

**Correspondence:** Diogo Sobreira Fernandes  
 Centro Hospitalar do Porto  
 Largo Prof. Abel Salazar  
 4099-001 Porto, Portugal  
 e-mail: sobreirafernandes@gmail.com

---